



On Artin Cokernel of The Group $D_n \times C_7$ When n is an Odd Number

Ass.Lec. Amer Khraja Abed Al-shypany
Al-muthanna Education, Ministry of Education

ASiraq@gmail.com

Received 23-3-2016,

Accepted 12-6-2016,

Published 13-10-2016

Abstract:

The group of all Z -valued generalized characters of G over the group of induced unit characters from all cyclic subgroups of G , $AC(G) = \bar{R}(G)/T(G)$ forms a finite abelian group, called Artin Cokernel of G . The problem of finding the cyclic decomposition of Artin cokernel $AC(D_n \times C_7)$ has been considered in this paper when n is an odd number, we find that if $n = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \dots \cdot p_m^{\alpha_m}$, where p_1, p_2, \dots, p_m are distinct primes and not equal to 2, then:

$$AC(D_n \times C_7) = \bigoplus_{i=1}^{2} C_2$$

$$= \bigoplus_{i=1}^2 AC(D_n) \oplus C_2$$

And we give the general form of Artin's characters table $Ar(D_n \times C_7)$ when n is an odd number.

Introduction

For a finite group G the finite abelian factor group $\bar{R}(G)/T(G)$ is called Artin cokernel of G and denoted by $AC(G)$ where $\bar{R}(G)$ denotes the abelian group generated by Z -valued characters of G under the operation of pointwise addition and $T(G)$ is a normal subgroup of $\bar{R}(G)$ which is generated by Artin's characters. Permutation characters induce from the principle characters of cyclic Subgroups. A well-known theorem which is due to Artin asserted that $T(G)$ has a finite index is,

i.e. $[\bar{R}(G):T(G)]$ is finite. The exponent of $AC(G)$ is called Artin exponent of G and denoted by $A(G)$. In 1968, Lam. T. Y [5] gave the definition of the group $AC(G)$ and studied $AC(C_n)$. In 1976, David. G [12] studied $A(G)$ of arbitrary characters of cyclic subgroups. In 1996, Knwabuez .K [11] studied $A(G)$ of p -groups. In 2000, H.R. Yassein [4] found $AC(G)$ for the group $\bigoplus_{i=1}^n C_p$. In 2002, k. Sekieguchi [12] studied the irreducible Artin characters of p -group and in the

Same year H. H. Abbass [10] found $\equiv^*(D_n)$.

In 2006, Abid. A. S [6] found $Ar(C_n)$ when C_n is the cyclic group of order n . In 2007, Mirza .R .N [9] found in her

1.Basic Concepts and Notations:

In this section, we recall some basic concepts, about matrix representation, characters and Artin character which will be used in later section.

Definition (1.1): [1]

A matrix representation of a group G is a homomorphism T of G into $GL(n, F)$, n is called the degree of matrix representation T . T is called a unit representation (principal) if $T(g)=1$, for all $g \in G$.

Definition (1.2):[2]

Let T be a matrix representation of a group G over the field F , the character χ of a matrix representation T is the mapping $\chi: G \rightarrow F$ defined by $\chi(g) = Tr(T(g))$ refers to the trace of the matrix $T(g)$ (the sum of the elements diagonal of $T(g)$). The degree of T is called the degree of χ .

Definition (1.3):[3]

Let H be acyclic subgroup of G and let ϕ be a class function on H . The induced class function on G is given by:

$$\phi'(g) = \frac{1}{|H|} \sum_{x \in G} \phi(xgx^{-1}), \forall g \in G$$

Where ϕ° is defined by:

$$\phi^\circ(h) = \begin{cases} \phi^\circ(h) & \text{if } h \in H \\ 0 & \text{if } h \notin H \end{cases}$$

Theorem (1.4):[4]

Let H be a cyclic subgroup of G and h_1, h_2, \dots, h_m are chosen representatives for Γ -conjugate classes, Then:

thesis Artin cokernel of the dihedral group. In this paper we find the general form of $Ar(D_n \times C_7)$ and we study $AC(D_n \times C_7)$ of the non abelian group $D_n \times C_7$ when n is an odd number.

$$\phi'(g) = \begin{cases} \frac{|C_G(g)|}{|C_H(g)|} \sum_{k=0}^m \phi^\circ(h_k) & \text{if } h_i \in H \cap CL(g) \\ 0 & \text{if } H \cap CL(g) = \emptyset \end{cases}$$

Definition (1.5):[5]

Let G be a finite group, all characters of G induced from the principal character of cyclic subgroups of G is called Artin characters of G .

Definition (1.6):[4]

Artin characters of the finite group can be displayed in a table called Artin characters table of G which is denoted by $Ar(G)$.

→ Proposition (1.7):[6]

The number of all distinct Artin characters on a group G is equal to the number of Γ -classes on G .

Definition (1.8):[1]

A rational valued character θ of G is a character whose values are in \mathbb{Z} , which is $\theta(g) \in \mathbb{Z}$, for all $g \in G$.

Definition (1.9):[6]

Let $T(G)$ be the subgroup of $\bar{R}(G)$ generated by Artin characters. $T(G)$ is a normal subgroup of $\bar{R}(G)$. Then the factor abelian group $\bar{R}(G)/T(G)$ is called Artin cokernel of G , denoted by $AC(G)$.

Proposition (1.10):[6]

$AC(G)$ is a finitely generated \mathbb{Z} -module.

Theorem [Artin] (1.11):[7]

Every rational valued character of G can be written as a linear combination of Artin characters with rational coefficient.

2. The Factor Group AC(G):

In this section, we use some concepts in linear algebra to study the factor group AC(G). We will give the general form of $Ar(D_n \times C_7)$ when n is an odd number. We shall study $Ac(G)$ dihedral group D_n and $\cong^*(D_n)$ when n is an odd number.

Definition (2.1):[5]

Let $T(G)$ be the subgroup of $\bar{R}(G)$ generated by Artin characters. $T(G)$ is a normal subgroup of $\bar{R}(G)$, then the factor abelian group $\bar{R}(G)/T(G)$ is called Artincokernel of G , denoted by $AC(G)$.

Definition (2.2): [8]

A_k -th determinant divisor of M is the greatest common divisor (g.c.d) of all the k -minors of M . This is denoted by $D_k(M)$.

Lemma (2.3): [8]

Let M, P and W be matrices with entries in the principal ideal domain R and p, W be invertible matrices, then: $D_k(P.M.W) = D_k(M)$ Modulo the group of units of R .

Theorem (2.4):[8]

Let M be an $k \times k$ matrix with entries in a principal ideal domain R , then there exists matrices P and W such that:

- 1 - P and W are invertible.
- 2 - $P M W = D$.
- 3 - D is a diagonal matrix.
- 4 - If we denote D_{jj} by d_j then there exists a natural number m ; $0 \leq m \leq k$ such that $j > m$ implies $d_j = 0$ and $j \leq m$ implies $d_j \neq 0$ and $1 \leq j \leq m$ implies $d_j | d_{j-1}$.

Definition (2.5):[8]

Let M be matrix with entries in a principal ideal domain R , equivalent to matrix $D = \text{diag} \{d_1, d_2, \dots, d_m, 0, 0,$

$\dots, 0\}$ such that $d_j | d_{j-1}$ for $1 \leq j < m$, we call D the invariant factor matrix of M and d_1, d_2, \dots, d_m the invariant factors of M .

Remark(2.6):

According to the Artin theorem (1.12) there exists an invertible matrix $M^{-1}(G)$ with entries in the set of rational numbers such that:

$\cong^*(G) = M^{-1}(G) \cdot Ar(G)$ and this implies, $M(G) = Ar(G) \cdot (\cong^*(G))^{-1}$ $M(G)$ is the matrix expressing the $T(G)$ basis in terms of the $\bar{R}(G)$ basis.

By theorem (2.5) there exists two matrices $P(G)$ and $W(G)$ with a determinant $\neq 1$ such that:

$$P(G) \cdot M(G) \cdot W(G) = \text{diag} \{d_1, d_2, \dots, d_l\} = D(G)$$

where $d_j = \pm D_j(G) | D_{j-1}(G)$ and l is the number of Γ -classes.

Theorem (2.7):[4]

$$AC(G) = \bigoplus_{i=1}^m Z \text{ where } d_j = \pm$$

$D_j(G) | D_{j-1}(G)$ where m is the number of all distinct Γ -classes.

Theorem(2.8):[9]

If n is an odd number such that $n = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \dots \cdot p_m^{\alpha_m}$, where p_1, p_2, \dots, p_m are distinct primes, then:

$$AC(D_n) = \bigoplus_{i=1}^{(\alpha_1 + 1) \cdot (\alpha_2 + 1) \cdot \dots \cdot (\alpha_m + 1) - 1} C_2$$

Proposition (2.9): [8]

The rational valued characters table of the cyclic group C_{p^s} of the rank $S+1$ where p is a prime number which is denoted by $(\cong^*(C_{p^s}))$, is given as follows:

Γ - classes	[1]	$[r^{P^{s-1}}]$	$[r^{P^{s-2}}]$	$[r^{P^{s-3}}]$...	$[r^{P^2}]$	$[r^P]$	[r]
θ_1	$P^{s-1}(P-1)$	$-P^{s-1}$	0	0	...	0	0	0
θ_2	$P^{s-2}(P-1)$	$P^{s-2}(P-1)$	$-P^{s-2}$	0	...	0	0	0
θ_3	$P^{s-3}(P-1)$	$P^{s-3}(P-1)$	$P^{s-3}(P-1)$	$-P^{s-3}$...	0	0	0
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
θ_{s-1}	$P(P-1)$	$P(P-1)$	$P(P-1)$	$P(P-1)$...	$P(P-1)$	$-P$	0
θ_s	$P-1$	$P-1$	$P-1$	$P-1$...	$P-1$	$P-1$	-1
θ_{s+1}	1	1	1	1	...	1	1	1

Table (2.1)

where its rank s+1 represents the number of all distinct Γ -classes.

Remark (2.10):[8]

If $n = p_1^{\alpha_1} \cdot p_1^{\alpha_2} \dots p_m^{\alpha_m}$, where p_1, p_2, \dots, p_m are distinct primes, then:

$$\equiv^*(C_n) = \equiv^*(C_{p_1^{\alpha_1}}) \otimes \equiv^*(C_{p_1^{\alpha_2}}) \otimes \dots \otimes \equiv^*(C_{p_m^{\alpha_m}}).$$

Definition (2.11):[7]

The dihedral group D_n is a certain non-abelian group of order $2n$. It is usually thought of as a group of transformations of the Euclidean plane of regular n -polygon consisting of rotations (about the origin) with the angle $2k\pi/n$, $k=0,1,2,\dots,n-1$ and

reflections (across lines through the origin). In general we can write it as :

$$D_n = \{ S^j r^k : 0 \leq k \leq n-1, 0 \leq j \leq 1 \}$$

which has the following properties :

$$r^n = 1, S^2 = 1, Sr^k S^{-1} = r^{-k}$$

Definition (2.12):

The group $D_n \times C_7$ is the direct product group $D_n \times C_7$, where C_7 is a cyclic group of order 7 consisting of elements $\{1, r', r^{2'}, r^{3'}, r^{4'}, r^{5'}, r^{6'}, r^{7'}\}$ with $(r')^7 = 1$. It is of order $14n$.

Theorem(2.13):[10]

The rational valued characters table of D_n when n is an odd number is given as follows:

$\equiv^*(C_n) =$		Γ - classes of C_n	[S]
	θ_1	$\equiv^*(C_n)$	0
	\vdots		\vdots
	θ_{s-1}		0
	θ_s	1 1 1 ... 1 1 1	1
	θ_{s+1}	1 1 1 ... 1 1 1	-1

Table (2.2)

Where S is the number of Γ - classes of C_n .

Theorem(2.14):

The rational valued characters table of the group $D_n \times C_7$ when n is an odd number is given as follows:

$$\equiv^*(D_n \times C_7) = \equiv^*(D_n) \otimes \equiv^*(C_7).$$

Theorem (2.15):[6]

The general form of Artin characters table of C_{p^s} when p is a prime number and s is positive integer is given by the lower Trianguler matrix:

- $M(C_{p_1^{\alpha_1}}) \otimes M(C_{p_2^{\alpha_2}}) \otimes \dots \otimes M(C_{p_m^{\alpha_m}}) \cdot M(C_n)$ is,
- 1- $(\alpha_1+1)(\alpha_2+1)\dots(\alpha_m+1) \times (\alpha_1+1)(\alpha_2+1)\dots(\alpha_m+1)$ square matrix.
 - 2- $P(C_n) = P(C_{p_1^{\alpha_1}}) \otimes P(C_{p_2^{\alpha_2}}) \otimes \dots \otimes P(C_{p_m^{\alpha_m}})$.
 - 3- $W(C_n) = W(C_{p_1^{\alpha_1}}) \otimes W(C_{p_2^{\alpha_2}}) \otimes \dots \otimes W(C_{p_m^{\alpha_m}})$.

3. The Main Results

In this section we give the general form of Artin characters table of the group $D_n \times C_7$ and the cyclic decomposition of the factor group $AC(D_n \times C_7)$ when n is an odd number.

Theorem(3.1):

The Artin characters table of the group $D_n \times C_7$ when n is an odd number is given as follows :

$Ar(D_n \times C_7) =$

Γ - classes	[1,1']	[1,r']	Γ - classes of $C_n \times C_7$					[S,1']	[S,r']
$ CL_\alpha $	1	1	2	2	2	n	n
$ C_{D_n \times C_7}(CL_\alpha) $	14n	14n	7n	7n	7n	14	14
$\Phi_{(1,1)}$	$2Ar(C_n) \otimes Ar(C_7)$							0	0
$\Phi_{(1,2)}$								⋮	⋮
⋮								⋮	⋮
$\Phi_{(l,1)}$								⋮	⋮
$\Phi_{(l,2)}$								0	0
$\Phi_{(l+1,1)}$	7n	0	0	0	7	0
$\Phi_{(l+1,2)}$	7n	0	0	0	0	7

Table(3.1)

where l is the number of Γ -classes of C_n and $C_7 = \langle r' \rangle$.

Proof:-By theorem(2.15)

$Ar(C_7) =$	Γ - classes	[1']	[r']
	$ CL_\alpha $	1	1
	$ C_7(CL_\alpha) $	7	7
	ϕ'_1	7	0
	ϕ'_2	1	1

Table (3.2)

Each cyclic subgroup of the group $D_n \times C_7$ is either a cyclic subgroup of $C_n \times C_7$ or (S,r') or $\langle (S,1') \rangle$. If H is a cyclic subgroup of $C_n \times C_7$, then :
 $H = H_i \times \langle 1' \rangle$ or $H_i \times \langle r' \rangle = H_i \times C_7$ for all $1 \leq i \leq l$ where l is the number of Γ -classes of C_n
 If $H = H_i \times \langle 1' \rangle$ and $x \in D_n \times C_7$

If $x \notin H$ then by theorem(1.4)
 $\Phi_{(1,i)}(x) = 0$ for all $0 \leq i \leq l$ [since $H \cap CL(x) = \phi$]
 If $x \in H$ then either $x = (1,1')$ or $\exists s, 0 < s < n$ such that $x = (r^s, 1')$
 If $x = (1,1')$, then :

$$\Phi_{(i,1)}(x) = \frac{|C_{D_n \times C_7}(x)|}{|C_H(x)|} \cdot \varphi'(x) \text{ [since } H \cap CL(x) = \{ (1, 1') \} \text{] ,}$$

where φ is the principle character

$$= \frac{14n}{|H_i| \cdot |<1'>|} \cdot 1 = \frac{14n}{|H_i|} = 2 \cdot \frac{n}{|H_i|} \cdot 1.7 = 2 \frac{|C_{C_n}(1)|}{|C_{H_i}(1)|} \cdot \varphi(1) \cdot \varphi'(1')$$

$$= 2 \cdot \varphi_i(1) \cdot \varphi'(1')$$

If $x=(r^s, 1')$ then

$$\Phi_{(i,1)}(x) = \frac{|C_{D_n \times C_7}(x)|}{|C_H(x)|} \cdot \sum_1^2 \varphi'(x) \text{ [since } H \cap CL(x) = \{(r^s, 1'), (r^{-s}, 1')\} \text{]}$$

$$= \frac{7n}{|H_{i \times <1'>}|} \cdot (1+1)$$

$$= \frac{7n}{|H_i|} \cdot 2$$

$$= 2 \cdot \frac{n}{|H_{i(r^s)}|} \cdot 1.7 = 2 \frac{|C_{C_n}(r^s)|}{|C_{H_i}(r^s)|} \cdot \varphi(r^s) \cdot \varphi'(1') = 2 \cdot \varphi_i(r^s) \cdot \varphi'_1(1')$$

If $H=H_i \times \langle r' \rangle = H_i \times C_7$

let $x \in D_n \times C_7$

if $x \notin H$ then

$\Phi_{(i,2)}(x) = 0$ for all $1 \leq i \leq l$ [since $H \cap CL(x) = \phi$]

If $x \in H$ then either $g=(1, 1')$ or $x=(1, r')$ or $\exists s, 0 < s < n$ such that $x= (r^s, r')$

If $x=(1, 1')$

$$\Phi_{(i,2)} = \frac{|C_{D_n \times C_7}(x)|}{|C_H(x)|} \cdot \varphi(x) \text{ [since } H \cap CL(x) = \{ (1, 1') \} \text{]}$$

$$= \frac{14}{|H_{i \times C_7}|} = \frac{14n}{2|H_i|} = \frac{7n}{|H_i|} = 2 \frac{|C_{C_n}(1)|}{|C_{H_i}(1)|} \cdot \varphi(1) = 7 \cdot \varphi_i(1) \cdot \varphi'_2(1)$$

If $x=(1, r')$ then

$$\Phi_{(i,2)}(x) = \frac{|C_{D_n \times C_7}(x)|}{|C_H(x)|} \cdot \varphi(x) \text{ [since } H \cap CL(x) = \{ (1, r') \} \text{]}$$

$$= \frac{14}{|H_{i \times C_7}|}$$

$$= \frac{14n}{2|H_i|} = \frac{7n}{|H_i|} = 7 \frac{|C_{C_n}(1)|}{|C_{H_i}(1)|} = 7 \cdot \varphi_i(1) \cdot \varphi'_2(r')$$

If $x=(r^s, r')$ then

$$\Phi_{(i,2)}(x) = \frac{|C_{D_n \times C_7}(x)|}{|C_H(x)|} \cdot \sum_1^2 \varphi'(x) \text{ [since } H \cap CL(x) = \{(r^s, r'), (r^{-s}, r')\} \text{]}$$

$$= \frac{7n}{|H_{i \times C_7}|} (1+1) = \frac{14n}{2|H_i|} = \frac{7n}{|H_i|} = 2 \frac{|C_{C_n}(r^s)|}{|C_{H_i}(r^s)|} \cdot \varphi(r^s) \cdot \varphi'_2(r') = 7 \cdot \varphi_i(r^s) \cdot \varphi'_2(r')$$

If $H = \langle (S, 1') \rangle = \{ (1, 1'), (S, 1') \}$ then:

$$\Phi_{(1+1,1)}((1, 1')) = \frac{|C_{D_n \times C_7(1, 1')}|}{|C_{H(S, 1')}|} \cdot \varphi(x) = \frac{14n}{2} = 7n$$

$$\Phi_{(1+1,1)}((1,1')) = \frac{|C_{D_n \times C_7(1,1')}|}{|C_{H(S,1')}|} \cdot \varphi(x) \text{ [since } H \cap CL(S, 1') = \{(S, 1')\} \text{]}$$

$$= \frac{14}{2} = 7$$

Otherwise

$$\Phi_{(1+1,1)}(x) = 0 \text{ for all } x \in D_n \times C_7, \text{ [since } x \notin H \text{]}$$

If $H = \langle (S,r') \rangle = \{ (1,1'), (S,r') \}$

$$\Phi_{(1+1,2)}((1,1')) = \frac{|C_{D_n \times C_7(1,1')}|}{|C_{H(1,1')}|} \cdot \varphi(1,1') \text{ [since } H \cap CL((1,1')) = \{(1,1')\} \text{]}$$

$$= \frac{14n}{2} \cdot 1 = 7n$$

$$\Phi_{(1+1,2)}((S,r')) = \frac{|C_{D_n \times C_7(S,r')}|}{|C_{H(S,r')}|} \cdot \varphi(S,r') = \frac{14}{2} \cdot 1 = 7$$

Otherwise $\Phi_{(1+1,2)}(x) = 0$ for all $x \in D_n \times C_7$ since $H \cap CL(x) = \phi$ ■

Proposition (3.2):

If $n = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \dots \cdot p_m^{\alpha_m}$ where p_1, p_2, \dots, p_m are distinct primes and $p_i \neq 2$ for all $1 \leq i \leq m$ and α_i any positive integers, then:

$$M(D_{n \times C_7}) = \begin{bmatrix} 2R(C_n) \times M(C_7) & 1 & 1 & 1 & 1 \\ 0 & 0 & \dots & \dots & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & \dots & \dots & 0 & 1 & 0 & 1 & 0 \\ 1 & 1 & \dots & \dots & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & \dots & \dots & 1 & 1 & 0 & 0 & 1 \end{bmatrix}$$

which is $2[(\alpha_1+1) \cdot (\alpha_2+1) \cdot \dots \cdot (\alpha_m+1) + 1] \times 2[(\alpha_1+1) \cdot (\alpha_2+1) \cdot \dots \cdot (\alpha_m+1) + 1]$ square matrix .

Proof:

By theorem(3.1) we obtain the Artin characters table $Ar(D_{n \times C_7})$ and from theorem(1.11) we find the rational valued characters table

$$\equiv^*(D_{n \times C_7}).$$

Thus by the definition of $M(G)$ we can find the 12×12 matrix $M(D_{n \times C_7})$:

$$\begin{aligned}
 & M(D_n \times C_7) = Ar(D_n \times C_7). (\\
 & \equiv *(D_n \times C_7))^{-1} = \\
 & \left[\begin{array}{cccccccccccc}
 2 & 2 & 2 & 2 & \dots & \dots & \dots & \dots & 1 & 1 & 1 & 1 \\
 0 & 2 & 0 & 2 & \dots & \dots & \dots & \dots & 1 & 0 & 1 & 0 \\
 0 & 0 & 2 & 2 & \dots & \dots & \dots & \dots & 1 & 1 & 1 & 1 \\
 0 & 0 & 0 & 2 & \dots & \dots & \dots & \dots & 1 & 0 & 1 & 0 \\
 & \vdots & \vdots & \vdots & \ddots & & & & \vdots & \vdots & \vdots & \vdots \\
 \vdots & \vdots & \vdots & \vdots & & \ddots & & & \vdots & \vdots & \vdots & \vdots \\
 \vdots & \vdots & \vdots & \vdots & & & \ddots & & \vdots & \vdots & \vdots & \vdots \\
 \vdots & \vdots & \vdots & \vdots & & & & \ddots & 1 & 0 & 1 & 0 \\
 0 & 0 & 0 & 0 & \dots & \dots & \dots & 0 & 1 & 1 & 1 & 1 \\
 0 & 0 & 0 & 0 & \dots & \dots & \dots & 0 & 1 & 0 & 1 & 0 \\
 1 & 1 & 1 & 1 & \dots & \dots & \dots & 1 & 1 & 1 & 0 & 0 \\
 1 & 1 & 1 & 1 & \dots & \dots & \dots & 1 & 1 & 0 & 0 & 1
 \end{array} \right] \\
 & = \left[\begin{array}{cccccccc}
 & & & & & 1 & 1 & 1 & 1 \\
 2R(C_n) \otimes M(C_7) & & & & & 1 & 0 & 1 & 0 \\
 & & & & & 1 & 1 & 1 & 1 \\
 & & & & & \vdots & \vdots & \vdots & \vdots \\
 0 & 0 & \dots & \dots & 0 & 1 & 1 & 1 & 1 \\
 0 & 0 & \dots & \dots & 0 & 1 & 0 & 1 & 0 \\
 1 & 1 & \dots & \dots & 1 & 1 & 1 & 0 & 0 \\
 1 & 1 & \dots & \dots & 1 & 1 & 0 & 0 & 1
 \end{array} \right]
 \end{aligned}$$

which is $2[(\alpha_1+1).(\alpha_2+1)... +1] \times 2[(\alpha_1+1).(\alpha_2+1)... +1]$ square matrix .

Proposition (3.3):

If $n = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \cdot \dots \cdot p_m^{\alpha_m}$ such that $g.c.d(P_i, P_j) = 1$ and $P_i \neq 2$ are prime numbers and α_i any positive integers, then:

$$P(D_{n \times C_7}) = \left[\begin{array}{cccccccccc}
 & & & & & & & & 0 & 0 \\
 & & & & & & & & 0 & 0 \\
 & & & & & & & & \vdots & \vdots \\
 & & & & & & & & 0 & 0 \\
 & & & & & & & & -1 & -1 \\
 & & & & & & & & 0 & 0 \\
 0 & 0 & 0 & \dots & 0 & 0 & 0 & & 1 & 0 \\
 0 & 0 & 0 & \dots & 0 & 0 & 0 & & 0 & 1
 \end{array} \right]$$

and

$$W(D_n \times C_7) = \begin{bmatrix} & & & & & & 0 & 0 & 0 & 0 \\ & & & & & & 0 & 0 & 0 & 0 \\ & & I_k & & & & \vdots & \vdots & \vdots & \vdots \\ & & & & & & \vdots & \vdots & \vdots & \vdots \\ & & & & & & 0 & 0 & 0 & 0 \\ & & & & & & 0 & 0 & 0 & 0 \\ -1 & -1 & \dots & \dots & -1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & \dots & \dots & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & 1 & \dots & \dots & 1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & \dots & \dots & 0 & 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$

Where $K = 2[(\alpha_1+1).(\alpha_2+1).(\alpha_3+1)...(\alpha_m+1) - 1] \times 2$
 $[(\alpha_1+1).(\alpha_2+1).(\alpha_3+1)...(\alpha_m+1) - 1]$

They are $2[(\alpha_1+1).(\alpha_2+1)...(\alpha_m+1)+1] \times 2$ $[(\alpha_1+1).(\alpha_2+1)...(\alpha_m+1) + 1]$ square matrix .

Proof :

By using theorem(2.5) and taking the form $M(D_n \times C_7)$ from proposition(3.2) and the above forms of $P(D_n \times C_7)$ and $W(D_n \times C_7)$ then we have

$$P(D_n \times C_7).M(D_n \times C_7).W(D_n \times C_7) = \begin{bmatrix} 2 & 0 & 0 & 0 & \dots & \dots & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & \dots & \dots & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & \dots & \dots & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & \dots & \dots & 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \dots & -2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & \dots & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & \dots & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & \dots & 0 & 0 & 0 & 1 \end{bmatrix}$$

$D(D_n \times C_7) = \text{diag}\{2, 2, 2, \dots, -2, 1, 1, 1\}$
 Which is $2[(\alpha_1+1).(\alpha_2+1)...(\alpha_m+1)+1] \times 2$ $[(\alpha_1+1).(\alpha_2+1)...(\alpha_m+1) + 1]$ square matrix .

Theorem (3.4):

If $n = p_1^{\alpha_1} . p_2^{\alpha_2} \dots p_m^{\alpha_m}$ where p_1, p_2, \dots, p_m are distinct prime numbers such that $P_i \neq 2$ and α_i any positive integers for all $i, 1 \leq i \leq m$, then the cyclic decomposition $AC(D_{n \times c_7})$ is :

$$AC(D_{n \times c_7}) = \bigoplus_{i=1}^{2((\alpha_1 + 1).(\alpha_2 + 1)...(\alpha_m + 1)) - 1} C_2$$

$$AC(D_{n \times c_7}) = \bigoplus_{i=1}^2 AC(D_n) \oplus C_2$$

Proof :

From proposition (3.3) we have
 $P(D_{n \times c_7}) . M(D_{n \times c_7}) . W(D_{n \times c_7}) = \text{diag}\{2, 2, 2, \dots, -2, 1, 1, 1\} = \{d_1, d_2, \dots, \dots, \dots\}$

$\{d_{2((\alpha_1+1),(\alpha_2+1),(\alpha_3+1),\dots,(\alpha_{m+1})) -1}, d_{2((\alpha_1+1),(\alpha_2+1),(\alpha_3+1),\dots,(\alpha_{m+1})) -1}$
 $,d_{2((\alpha_1+1),(\alpha_2+1),(\alpha_3+1),\dots,(\alpha_{m+1}))},$
 $d_{2((\alpha_1+1),(\alpha_2+1),(\alpha_3+1),\dots,(\alpha_{m+1})) +1}d_{2((\alpha_1+1),(\alpha_2+1),(\alpha_3+1),\dots,(\alpha_{m+1})) +2} \}$.
 By theorem (2.8) we get

$$2((\alpha_1 + 1).(\alpha_2 + 1)\dots(\alpha_m + 1)) -1$$

$$AC(D_{n \times c_7}) = \bigoplus_{i=1}^2 C_{d_i}$$

$$= \bigoplus_{i=1}^{2((\alpha_1 + 1).(\alpha_2 + 1)\dots(\alpha_m + 1)) -1} C_2$$

From theorem(2.9) we have :

$$AC(D_{n \times c_7}) = \bigoplus_{i=1}^2 AC(D_n) \oplus C_2$$

Example (3.6):

To find the cyclic decomposition of the groups $AC(D_{12167 \times C_7})$, $AC(D_{11692487 \times C_7})$ and $AC(D_{222157253 \times C_7})$.

We can use above theorem :

$$1-AC(D_{12167 \times C_7}) = AC(23^3 \times C_7) = \bigoplus_{i=1}^{2(3+1)-1} C_2 = \bigoplus_{i=1}^7 C_2 = \bigoplus_{i=1}^2 AC(D_{23^3}) \oplus C_2 .$$

$$2-AC(D_{11692487 \times C_7}) = AC(D_{23^3 \cdot 31^2 \times C_7}) = \bigoplus_{i=1}^{2((3+1).(2+1))-1} C_2 = \bigoplus_{i=1}^{15} C_2$$

$$= \bigoplus_{i=1}^2 AC(D_{23^3 \cdot 31^2}) \oplus C_2$$

$$3-AC(D_{222157253 \times C_7}) = AC(D_{23^3 \cdot 31^2 \cdot 19 \times C_7}) = \bigoplus_{i=1}^{2((3+1).(2+1).(1+1))-1} C_2$$

$$= \bigoplus_{i=1}^{47} C_2 = \bigoplus_{i=1}^2 AC(D_{23^3 \cdot 31^2 \cdot 19}) \oplus C_2$$

References

[1] Culirits . C and.Reiner . I, "Methods of Representation Theory with Appcation to Finite Groups and Order", John wily & sons, New york, 1981.

[2] A. M, Basheer "Representation Theory of Finite Groups" ,AIMS, South Africa , 2006.
 [3] J Moori, "Finite Groups and Representation Theory" ,University of Kawzulu–Natal , 2006 .

- [4] H. R Yassien , "On ArtinCokernel of Finite Group", M.Sc. Thesis, Babylon niversity, 2000.
- [5] T .Y Lam, "Artin Exponent of Finite Groups", Columbia University, New York, 1968.
- [6] A. S, Abid. "Artin's Characters Table of Dihedral Group for Odd Number", MSc.Thesis, university of kufa, 2006.
- [7] J. P Serre, "Linear Representation of Finite Groups", Springer- Verlage, 1977.
- [8] M . S Kirdar .M . S, "The Factor Group of The Z- Valued Class Function Modulo the Group of The Generalized Characters ", Ph. D. Thesis, University of Birmingham, 1982.
- [9] Mirza . R. N, "On ArtinCokernel of Dihedral Group Dn When n is An Odd Number", M.Sc. thesis, University of Kufa , 2007.
- [10] H .H Abass, "On Rational of Finite Group D_n when n is an odd", Journal Babylon University, vol7, No-3, 2002.
- [11] Knwabusz . K, "Some Definitions of Artin's Exponent of Finite Group", USA.National foundation Math.GR.1996.
- [12] David .G, "Artin Exponent of Arbitrary Characters of Cyclic Subgroups", Journal of Algebra, 61, pp.58-76, 1976.

حول النواة المشارك - ارتن للزمرة $D_n \times C_7$ عندما n عدد فردي

م.م. عامر خريجة عبد الشيباني
مديرية تربية المثني/ وزارة التربية

الخلاصة

أن زمرة كل الشواخص العمومية ذات القيم الصحيحة للزمرة G على زمرة الشواخص المحتثة من الشواخص الأحادية للزمرة الجزئية الدائرية $AC(G) = \bar{R}(G)/T(G)$ تكون زمرة ابيلية منتهية وتسمى النواة المشارك- ارتن للزمرة G . إن مسألة إيجاد التجزئة الدائرية للزمرة القسمة $AC(G)$ تم اعتبارها في هذا البحث للزمرة $D_n \times C_7$ عندما n عدد فرد، وجدنا إذا كانت $n = p_1^{\alpha_1} \cdot p_1^{\alpha_2} \dots p_m^{\alpha_m}$ وإن p_1, p_2, \dots, p_m أعداد أولية مختلفة لا تساوي 2 فإن:

$$2((\alpha_1 + 1) \cdot (\alpha_2 + 1) \dots (\alpha_m + 1)) - 1$$

$$AC(D_n \times C_7) = \bigoplus_{i=1}^2 C_2$$

$$= \bigoplus_{i=1}^2 AC(D_n) \oplus C_2$$

وكذلك وجدنا الصيغة العامة لجدول شواخص ارتن $Ar(D_n \times C_7)$ عندما يكون n عدد فردي.